

Bcl-2 (phospho-S70) polyclonal antibody

Catalog: BCP00271

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Bcl-2 is one among many key regulators of apoptosis which are essential for proper development, tissue homeostasis and protection against foreign pathogens. Human Bcl-2 is a 26 kDa, membrane-associated, anti-apoptotic oncoprotein that can promote cell survival through protein-protein interactions with other Bcl-2 related family members, such as the death suppressors Bcl-xL, Mcl-1, Bcl-w and A1, or the death agonists Bax, Bak, Bik, Bad and BID. The anti-apoptotic function of Bcl-2 can also be regulated through proteolytic processing and phosphorylation. Bcl-2 may promote cell survival by interfering with the activation of the cytochrome c/Apaf-1 pathway through stabilization of the mitochondrial membrane.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 26 kDa

Swiss-Prot:

P10415

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IP: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-Bcl-2 (S70) polyclonal antibody detects endogenous levels of Bcl-2 protein only when phosphorylated at Ser70.

DATA:




Western blot (WB) analysis of Bcl-2 (phospho-S70) polyclonal antibody at 1:500 dilution

Lane1: HEK293T whole cell lysate(40ug)

Lane2: HEK293T treated with UV for 1h whole cell lysate(40ug)

Lane3: HEK293T treated with UV for 6h whole cell lysate(40ug)

Lane4: HEK293T treated with UV for 24h whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of p-Bcl2 (S70) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.