

Catenin- β (I35) polyclonal antibody

Catalog: BCP00346

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

β -Catenin is a key downstream effector in the Wnt signaling pathway. It is implicated in two major biological processes in vertebrates: early embryonic development and tumorigenesis. CK1 phosphorylates β -catenin at Ser45. This phosphorylation event primes β -catenin for subsequent phosphorylation by GSK-3 β . GSK-3 β destabilizes β -catenin by phosphorylating it at Ser33, Ser37, and Thr41. Mutations at these sites result in the stabilization of β -catenin protein levels and have been found in many tumor cell lines.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 92 kDa

Swiss-Prot:

P35222

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Catenin- β (I35) polyclonal antibody detects endogenous levels of Catenin- β protein.

DATA:




Western blot (WB) analysis of Catenin- β (I35) pAb at 1:500 dilution

Lane1:Hela whole cell lysate(40ug)

Lane2:The Uterus tissue lysate of Rat(40ug)

Lane3:The Embryo tissue lysate of Mouse(40ug)



Note:

For research use only, not for use in diagnostic procedure.