

## CD4 (10H8) monoclonal antibody

Catalog: BCP00445

Host: Mouse

Reactivity: Human,Mouse,Rat

### BackGround:

Cluster of Differentiation 4 (CD4) is a glycoprotein composed of an amino-terminal extracellular domain (four domains: D1-D4 with Ig-like structures), a trans-membrane part and a short cytoplasmic tail. CD4 is expressed on the surface of T helper cells, regulatory T cells, monocytes, macrophages and dendritic cells, and plays an important role in the development and activation of T cells. On T cells, CD4 is the co-receptor for the T cell receptor (TCR), and these two distinct structures recognize the Antigen-Major Histocompatibility Complex (MHC). Specifically, the D1 domain of CD4 interacts with the  $\beta$ 2-domain of the MHC class II molecule. CD4 ensures specificity of the TCR-antigen interaction, prolongs the contact between the T cell and the antigen presenting cell and recruits the tyrosine kinase Lck, which is essential for T cell activation.

### Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 51 kDa

### Swiss-Prot:

P01730

### Purification&Purity:

The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immuno-

gen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

CD4 ( 10H8) monoclonal antibody detects endogenous levels of CD4 protein.

### DATA:



Western blot (WB) analysis of CD4 (10H8) mAb at 1:1000 dilution

Lane1:Jurkat whole cell lysate(30ug)

Lane2:THP-1 whole cell lysate(30ug)

### Note:

For research use only, not for use in diagnostic procedure.