COMPLEX

CD48 polyclonal antibody

Catalog:	BCP00449	Host:	Rabbit	Reactivity: Human,Rat
BackGrou	nd•			munogen and the purity is $> 95\%$ (by SDS-PAGE).
CD48 is a glycosylphosphatidylinositol (GPI) -anchored				Applications:
membrane protein of the signaling lymphocyte activation				WB: 1:1000~1:2000
molecule (SLAM) family, also known as SLAMF2 and				Storage&Stability:
BLAST-1. It is constitutively expressed on most hemato-				
poietic cells (not on neutrophils and a subset of long-term				Store at $4 ^{\circ}$ C short term. Aliquot and store at $-20 ^{\circ}$ C long term. Avoid fracts there evalues
hematopoietic stem cells in mice) and can be upregulated				term. Avoid freeze-thaw cycles.
under certain conditions like infection. Interaction with its			1 0	Specificity:
low affinity ligand CD2 promotes adhesion and TCR				CD48 polyclonal antibody detects endogenous levels of
signaling. Interaction with the high affinity ligand CD244				CD48 protein.
(2B4) regulates natural killer (NK) and CD8 T cell acti-			-	DATA:
vation and cytolytic function.				
Product:				Western blot (WB) analysis of CD48 polyclonal antibody at 1:1000 di-
			dinne enide	lution
Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2. Molecular Weight: ~ 48 kDa			odium azide,	Lane1:LOVO whole cell lysate(30ug)
				Lane2:L02 cell membrane lysate(30ug)
				Lane3:MCF-7 whole cell lysate(30ug)
				Lane4:PMVEC whole cell lysate(20ug)
Swiss-Prot:				Note:
P09326				For research use only, not for use in diagnostic procedure.
Purification&Purity:				

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-