Cdk1/Cdc2 (phospho-T161) polyclonal antibody

Catalog: BCP00479

Host: Rabbit

Reactivity:

Human, Mouse, Rat

BackGround:

Cdc2. evolutionarily conserved an serine/threonine-specific protein kinase, is essential in the cell cycle transition from G2 to M phase. Cdc2 is regulated by association with B-type cyclins and by reversible phosophorylation. Cyclin B binding facilitates the phosphorylation of Cdc2 p34 on three regulatory sites: threonine 14, tyrosine 15, and threonine 161. In higher eukaryotes, Cdc2 is negatively regulated by phosphorylation of two residues located in the ATP-binding site, Thr 14 and Tyr 15. Cdc2 is positively regulated by the cyclin-dependent phosphorylation of Thr 161. Both phosphorylation and de- phosphorylation at Thr 161 are required for progression through the cell cycle.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 34 kDa

Swiss-Prot:

P06493

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

Storage&Stability:

Store at $4 \ \mathbb{C}$ short term. Aliquot and store at $-20 \ \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

p-Cdk1/Cdc2 (T161) polyclonal antibody detects endog-

enous levels of p-Cdk1/Cdc2 protein only when phosphorylated at Thr161.

DATA:



Western blot (WB) analysis of Cdk1/Cdc2 (phospho-T161) polyclonal antibody at 1:500 dilution

Lane1:PC12 whole cell lysate(40ug) Lane2:3T3-L1 whole cell lysate(40ug) Lane3:HEK293T whole cell lysate(40ug) Lane4:MCF-7 whole cell lysate(40ug) Lane5:PC3 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of Cdk1/Cdc2 (phospho-T161) polyclonal antibody in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.