

c-IAP2 polyclonal antibody

Catalog: BCP00498

Host: Rabbit

Reactivity: Rat, Mouse

BackGround:

The inhibitor of apoptosis protein (IAP) family consists of an evolutionarily conserved group of apoptosis inhibitors containing a conserved 70 amino acid BIR (baculovirus inhibitor repeat) domain. Human members of this family include c-IAP1, c-IAP2, XIAP, survivin, livin, and NAIP. Overexpression of IAP family members, particularly survivin and livin, in cancer cell lines and primary tumors suggests an important role for these proteins in cancer progression. In general, the IAP proteins function through direct interactions to inhibit the activity of several caspases, including caspase-3, caspase-7, and caspase-9. In addition, binding of IAP family members to the mitochondrial protein Smac blocks their interaction with caspase-9, thereby allowing the processing and activation of the caspase.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~ 68 kDa

Swiss-Prot:

Q13489

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:


WB: 1:1000~1:2000

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

c-IAP2 polyclonal antibody detects endogenous levels of c-IAP2 protein.

DATA:

Western blot (WB) analysis of c-IAP2 polyclonal antibody at 1:1000 dilution

Lane1: The Heart tissue lysate of Rat(40ug)

Lane2: The Liver tissue lysate of Mouse(40ug)

Note:

For research use only, not for use in diagnostic procedure.