

CNOT7 (N19) polyclonal antibody

Catalog: BCP00537

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

CNOT7 (CCR4-NOT transcription complex, subunit 7), also known as CAF1 (CCR4-associated factor 1), hCAF-1 or BTG1-binding factor 1, is a member of the CAF1 family. Localizing to the nucleus, CNOT7 is ubiquitously expressed and is believed to function as a transcription factor, playing a role in a wide variety of processes. CNOT7 functions as a component of the evolutionarily conserved CCR4-NOT complex, a multi-subunit complex that participates in transcription as well as mRNA degradation. CNOT7 and other subunits of the CCR4-NOT complex play a role in the regulation of nuclear hormone receptor activities. CNOT7 directly binds to and interacts with RXR β , TOB1, TOB2, BTG1, BTG2 and BTG3. In addition, CNOT7 knockout mice are sterile and show an increase in bone mass, suggesting an important role for CNOT7 in spermatogenesis and as a suppressor of bone mass and BMP (bone morphogenetic protein) actions in osteoblasts.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 32 kDa

Swiss-Prot:

Q9UIV1

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

CNOT7 (N19) polyclonal antibody detects endogenous levels of CNOT7 protein.

DATA:



Western blot (WB) analysis of CNOT7 (N19) Antibody at 1:1000 dilution

Lane1:Myla2059 whole cell lysate(40ug)

Lane2:HEK293T whole cell lysate(40ug)

Lane3:Hela whole cell lysate(40ug)

Lane4:PC12 whole cell lysate(40ug)

Lane5:3T3-L1 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.