

Cyclin D1 (L283) polyclonal antibody

Catalog: BCP00603

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Considerable effort directed towards the identification of G1 cyclins has led to the isolation of cyclin D, cyclin C and cyclin E. Of these, cyclin D corresponds to a putative human oncogene, designated PRAD1, which maps at the site of the Bcl1 rearrangement in certain lymphomas and leukemias. Two additional human type D cyclins, as well as their mouse homologs, have been identified. Evidence has established that members of the cyclin D family function to regulate phosphorylation of the retinoblastoma gene product, thereby activating E2F transcription factors.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 36 kDa

Swiss-Prot:

P24385

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Cyclin D1 (L283) polyclonal antibody detects endogenous levels of Cyclin D1 protein.

DATA:




Western blot (WB) analysis of Cyclin D1 (L283) pAb at 1:2000 dilution

Lane1:HepG2 whole cell lysate(20ug)

Lane2:L02 whole cell lysate(20ug)

Lane3:C6 whole cell lysate(20ug)

Lane4:3T3-L1 whole cell lysate(20ug)



Immunohistochemistry (IHC) analyzes of Cyclin D1 (L283) pAb in paraffin-embedded human breast carcinoma tissue at 1:50.

Note:

For research use only, not for use in diagnostic procedure.