

CYP27A1 (E131) polyclonal antibody

Catalog: BCP00612

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

CYP27A1 participates in the degradation of cholesterol to bile acids in both the classic and acidic pathways.[1] It is the initiating enzyme in the acidic pathway to bile acids, yielding oxysterols by introducing a hydroxyl group to the carbon at the 27 position in cholesterol. In the acidic pathway, it produces 27-hydroxycholesterol from cholesterol whereas in the classic or neutral pathway, it produces 3 β -hydroxy-5-cholestenoic acid. It is also involved in the metabolism of vitamin D3. While CYP27A1 is present in many different tissues, its function in these tissues is largely uncharacterized. In macrophages, 27-hydroxycholesterol generated by this enzyme may be helpful against the production of inflammatory factors associated with cardiovascular disease.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 60 kDa

Swiss-Prot:

Q02318

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


IF: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

CYP27A1 (E131) polyclonal antibody detects endogenous levels of CYP27A1 protein

DATA:

Western blot (WB) analysis of CYP27A1 (E131) pAb at 1:500 dilution


Lane1: LO2 whole cell lysate(40ug)

Lane2: HepG2 whole cell lysate(40ug)

Lane3: A549 whole cell lysate(40ug)

Lane4: PC12 whole cell lysate(40ug)

Lane5: AML-12 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of CYP27A1 (E131) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.