DAPK3 (phospho-T265) polyclonal antibody

Catalog: BCP00639

Host: R

Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Apoptosis is mediated by death domain containing adapter molecules and a caspase family of proteases. Certainserine/threonine protein kinases, such as ASK1 and RIP, are mediators of apoptosis. A novel serine/threonine kinase that mediates apoptosis was recently identified and designated ZIP kinase. ZIP kinase contains an N terminal kinase domain and a C terminal leucine zipper structure and binds to ATF4 that is a member of ATF/CREBfamily. ZIP kinase has high sequence homology to DAP kinase (death associated protein kinase), which is a mediator of apoptosis induced by gamma interferon. Overexpression of ZIP kinase induces apoptosis. ZIP and DAP kinases represent a novel kinase family, which mediates apoptosis through their catalytic activities. The messenger RNA was ubiquitously expressed in various tissues.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 52 kDa

Swiss-Prot:

O43293

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

DAPK3 (phospho-T265) polyclonal antibody detects endogenous levels of DAPK3 protein when phosphorylated at Thr265.

DATA:



Western blot (WB) analysis of p-DAPK3 (T265) pAb at 1:500 dilution Lane1:Hela whole cell lysate(40 µg) Lane2:SK-OVCAR3 whole cell lysate(40 µg) Lane4:3T3-L1 whole cell lysate(40 µg) Lane5:PC12 whole cell lysate(40 µg)

Note:

For research use only, not for use in diagnostic procedure.