

Ephrin-A3 (E196) polyclonal antibody

Catalog: BCP00729

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The Eph subfamily represents the largest group of receptor protein kinases identified to date. There is increasing evidence that they are involved in central nervous system function and in development. Ligands for Eph receptors include ephrin-A1 (LERK-1/B61), identified as a ligand for the EphA2 (Eck) receptor; ephrin-A2 (ELF-1), identified as a ligand for the EphA3 and EphA4 (Sek) receptors; ephrin-A3 (LERK-3), identified as a ligand for EphA5 (Ehk1) and EphA3 (Hek) receptors; ephrin-A4 (LERK-4), identified as a ligand for the EphA3 receptor; ephrin-A5 (AL-1), identified as a ligand for EphA5 (REK7); ephrin-B1 (LERK-2), identified as a ligand for the EphB1 (Elk) and EphB2 (Cek5) receptors; ephrin-B2 (LERK-5), identified as a ligand for the EphB1, EphB3 (Cek10) and EphB2 receptors; and ephrin-B3 (LERK-8), identified as a ligand for EphB1.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 38 kDa

Swiss-Prot:

P52797

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Ephrin-A3 (E196) polyclonal antibody detects endogenous levels of Ephrin-A3 protein.

DATA:




Western blot (WB) analysis of Ephrin-A3 (E196)pAb at 1:500 dilution

Lane1:Hct116 whole cell lysate

Lane2:SK-OVCAR3 whole cell lysate

Lane3:The Brain tissue lysate of Mouse

Lane4:The Brain tissue lysate of Rat



Immunohistochemistry (IHC) analyzes of Ephrin-A3 (E196) pAb in paraffin-embedded human brain tissue.

Note:

For research use only, not for use in diagnostic procedure.