

PUMA (R169) polyclonal antibody

Catalog: BCP01394

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

PUMA (Bcl-2 binding component 3, JFY1, PUMA/JFY1) is a mitochondrial pro-apoptotic Bcl-2 homology domain (BH3)-only protein that induces rapid apoptosis through a Bax- and mitochondria-dependent pathway. The PUMA gene encodes four proteins originating from different splice variants of the same transcript: PUMA α , β , γ and δ . Both PUMA α and PUMA β contain a BH3 domain, while PUMA γ and PUMA δ lack this domain. The BH3 domain is essential for binding of PUMA α and PUMA β to Bcl-2 or Bcl-xL. PUMA is an initiator of γ -radiation apoptosis and glucocorticoid-induced apoptosis in lymphoid cells in vivo. Bcl-2 family members generally regulate apoptosis and transmit death signals to mitochondria. Members of this family include both pro- and anti-apoptotic proteins that share homologous sequences known as Bcl-2 homology domains (BH1-4). The BH3 proteins, BID, NOXA, PUMA, NBK, Bim and Bad, are all pro-apoptotic and share sequence homology within the amphipathic α -helical BH3 region.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 15, 20, 28 kDa

Swiss-Prot:

Q9BXH1

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PUMA (R169) polyclonal antibody detects endogenous levels of PUMA protein.

DATA:



Western blot (WB) analysis of PUMA (R169) polyclonal antibody at 1:500 dilution

Lane1:L02 whole cell lysate(40ug)

Lane2:HepG2 whole cell lysate(40ug)

Lane3:MCF-7 whole cell lysate(40ug)

Lane4:PC12 whole cell lysate(40ug)

Lane5:BV2 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.