

## PXR polyclonal antibody

Catalog: BCP01395

Host: Rabbit

Reactivity: Human,Pig

### BackGround:

Steroid hormones function as signaling molecules by diffusing into cells and interacting with specific intracellular receptors to regulate gene expression. This superfamily of receptors includes both steroid and nonsteroid receptors. Like many nonsteroid hormone receptors, PXR (pregnane X receptor) binds as a heterodimer with RXR to a DNA sequence typical of a nonsteroid hormone receptor; however, PXR is activated by several steroids, such as naturally occurring pregnanes and synthetic glucocorticoids and anti-glucocorticoids. PXR exists as two alternatively spliced isoforms, PXR.1 and PXR.2. PXR is thought to define a novel steroid hormone signaling pathway that may account for some of the effects of synthetic glucocorticoids and antiglucocorticoids that are not mediated through the classical glucocorticoid receptor signaling pathway.

### Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

~ 52 kDa

### Swiss-Prot:

O75469

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:1000~1:2000


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

PXR polyclonal antibody detects endogenous levels of PXR protein.

### DATA:



Western blot (WB) analysis of PXR polyclonal antibody at 1:2000 dilution

Lane1:HepG2 whole cell lysate(40ug)

Lane2:THP-1 whole cell lysate(40ug)

Lane3:The Liver tissue lysate of Pig(40ug)

### Note:

For research use only, not for use in diagnostic procedure.