

Rsk-1 (T353) polyclonal antibody

Catalog: BCP01482 Host: Rabbit Reactivity: Human, Rat

BackGround:

The family of ribosomal S6 kinases (Rsks), designated Rsk-1 (or MAPKAP kinase-1), Rsk-2 and Rsk-3, are intracellular serine/threonine kinases that are important signaling intermediates in response to a broad range of ligand activated receptor tyrosine kinases. A unique feature common to the members of the Rsk family is that each possesses two non-identical complete kinase catalytic domains. An additional Rsk protein, Rsk-4, shows a high level of homology to the three previously isolated members of the human Rsk family. Rsk-4 is most abundantly expressed in brain and kidney and plays a role in normal neuronal development.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 85 kDa

Swiss-Prot:

Q15418

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:2000~1:5000 IF: 1:50~1:200 IP: 1:50~1:200


Storage&Stability:

Store at $4 \,\mathrm{C}$ short term. Aliquot and store at $-20 \,\mathrm{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

Rsk-1 (T353) polyclonal antibody detects endogenous levels of Rsk-1 protein.

DATA:




Western blot (WB) analysis of Rsk-1 (T353) pAb at 1:2000 dilution

Lane1:C6 whole cell lysate(40ug)


Lane2:HCT116 whole cell lysate(40ug)

Lane3:Jurkat whole cell lysate(40ug)

Lane4:H1792 whole cell lysate(40ug)



Immunoprecipitation of HCT116 cell lysates using Rsk-1 pAb (Sepharose Bead Conjugate)#BD0048 (lane 3 and lane 4) and Nonspecific IgG Control (Sepharose Bead Conjugate)#BD0048 (lane 1 and lane 2). Lane 1 is 30% input. The western blot was probed using Rsk-1 pAb.



Immunofluorescence analysis of HCT116 cells using Rsk-1 antibody at dilution of 1:50.

Note:

For research use only, not for use in diagnostic procedure.