

Smad2/3 (phospho-T8) polyclonal antibody

Catalog: BCP01535

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

Smad proteins, the mammalian homologs of the Drosophila mothers against decapentaplegic (Mad), have been implicated as downstream effectors of GF β /BMP signaling. Smad1 (also designated Madr1 or JV4-1) and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF β and activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to activin/TGF β signaling by interfering with TGF β -mediated phosphorylation of other Smad proteins.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 55, 60 kDa

Swiss-Prot:

P84022/Q15796

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-Smad2/3 (T8) polyclonal antibody detects endogenous levels of Smad2/3 protein only when phosphorylated at Thr8.

DATA:



Western blot (WB) analysis of p-Smad2/3 (T8) pAb at 1:500 dilution

Lane1:HEK293T whole cell lysate(40 µg)

Lane2:Hela whole cell lysate(40 µg)

Lane3:The Lung tissue lysate of Mouse(40 µg)

Lane4:SGC7901 whole cell lysate(40 µg)

Lane5:H9C2 whole cell lysate(40 µg)

Note:

For research use only, not for use in diagnostic procedure.