

Smad3 (phospho-S213) polyclonal antibody

Catalog: BCP01539

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

Smad proteins, the mammalian homologs of the Drosophila mothers against decapentaplegic (Mad), have been implicated as downstream effectors of $\text{GF}\beta$ /BMP signaling. Smad1 (also designated Madr1 or JV4-1) and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in $\text{TGF}\beta$ and activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to activin/ $\text{TGF}\beta$ signaling by interfering with $\text{TGF}\beta$ -mediated phosphorylation of other Smad proteins.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 48, 55 kDa

Swiss-Prot:

P84022

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-Smad3 (S213) polyclonal antibody detects endogenous levels of Smad3 protein only when phosphorylated at Ser213.

DATA:



Western blot (WB) analysis of p-Smad3 (S213) pAb at 1:500 dilution

Lane1:Hela whole cell lysate(40ug)

Lane2:HEK293T whole cell lysate(40ug)

Lane3:Panc1 whole cell lysate(20ug)

Lane4:LOVO whole cell lysate(40ug)

Lane5:C6 whole cell lysate(40ug)

Lane6:AML-12 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.