

**TGFβ2 (L388) polyclonal antibody**

Catalog: BCP01637

Host: Rabbit

Reactivity: Human, Mouse, Rat

**BackGround:**

Transforming growth factor betas (TGFβs) were originally discovered due to their ability to promote anchorage-independent growth of rat NRK fibroblasts in the presence of TGFα. It is now realized that TGFβs mediate many cell-cell interactions that occur during embryonic development. Three TGFβs have been identified in mammals. TGFβ1, TGFβ2 and TGFβ3 are each synthesized as precursor proteins that are very similar in that each is cleaved to yield a 112 amino acid polypeptide that remains associated with the latent portion of the molecules. Biologically active TGFβ requires dimerization of the monomers (usually homodimers) and release of the latent peptide portion. Overall, the mature region of the TGFβ3 protein has approximately 80% identity to the mature region of both TGFβ1 and TGFβ2. However, the NH2 terminals or precursor regions of their molecules share only 27% sequence identity.

**Product:**

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

**Molecular Weight:**

~ 48; 55 kDa

**Swiss-Prot:**

P61812

**Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

**Applications:**

WB: 1:500~1:1000


IHC: 1:50~1:200

**Storage&Stability:**

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

**Specificity:**

TGF β2 (L388) polyclonal antibody detects endogenous levels of TGF β2 protein.


**DATA:**

Western blot (WB) analysis of TGFβ2 (L388) pAb at 1:500 dilution

Lane1:HEK293T whole cell lysate(40ug)

Lane2:A375 whole cell lysate(40ug)

Lane3:MCF-7 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of TGFβ2 (L388) pAb in paraffin-embedded human lung carcinoma tissue at 1:100.

**Note:**

For research use only, not for use in diagnostic procedure.