Tie-2 (phospho-Y1108) polyclonal antibody

Catalog: BCP01644 Host: Rabbit Reactivity: Human,Mouse,Rat

BackGround:

Receptor tyrosine kinases play key roles in signal transduction across cell surfaces in biological systems, including the vascular system. These receptors comprise a large and diverse family of catalytically related proteins that, on the basis of sequence and structural similarities, can be divided into several different evolutionary subfamilies. The cloning and characterization of Tie-1 (also designated Tie), a novel human endothelial cell surface receptor tyrosine kinase, has been reported. The extracellular domain of the predicted Tie-1 protein product has an unusual multidomain structure consisting of a cluster of three epidermal growth factor homology motifs localized between two immunoglobulin-like loops, which are followed by three fibronectin type III repeats next to the transmembrane region. An additional member of this family has been identified as Tie-2 (also designated Tek). Tie-1 and Tie-2 have been shown to be encoded by distinct genes and to represent members of a new class of receptor tyrosine kinases.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 160 kDa

Swiss-Prot:

Q02763

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:2000

Storage&Stability:

Store at 4℃ short term. Aliquot and store at -20℃ long term. Avoid freeze-thaw cycles.

Specificity:

p-Tie-2 (Y1108) polyclonal antibody detects endogenous levels of Tie-2 protein when phosphorylated at Tyr1108.

DATA:



Western blot (WB) analysis of p-Tie-2 (Y1108) pAb at 1:2000 dilution

Lane1:A549 whole cell lysate(40ug)

Lane2:A375 whole cell lysate(40ug)

Lane3:SGC7901 whole cell lysate(40ug)

Lane4:H9C2 whole cell lysate(40ug)

Lane5:AML-12 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Complex biotech, co. Ltd.
Add：No 9, weidi road Qixia District Nanjing, 210046, P. R. China.