

XIAP polyclonal antibody

Catalog: BCP01725

Host: Rabbit

Reactivity: Human,Rat,Mouse

BackGround:

The inhibitor of apoptosis protein (IAP) family consists of an evolutionarily conserved group of apoptosis inhibitors containing a conserved 70 amino acid BIR (baculovirus inhibitor repeat) domain. Human members of this family include c-IAP1, c-IAP2, XIAP, survivin, livin, and NAIP. Overexpression of IAP family members, particularly survivin and livin, in cancer cell lines and primary tumors suggests an important role for these proteins in cancer progression. In general, the IAP proteins function through direct interactions to inhibit the activity of several caspases, including caspase-3, caspase-7, and caspase-9. In addition, binding of IAP family members to the mitochondrial protein Smac blocks their interaction with caspase-9, thereby allowing the processing and activation of the caspase.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~ 60 kDa

Swiss-Prot:

P98170

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 96% (by SDS-PAGE).

Applications:

WB: 1:5000~1:10000

IP:1:2000~1:8000

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

Specificity:

XIAP polyclonal antibody detects endogenous levels of XIAP protein.

DATA:




Western blot (WB) analysis of XIAP polyclonal antibody at 1:5000 dilution

Lane1:H9C2 whole cell lysate(40ug)

Lane2:AML-12 whole cell lysate(40ug)

Lane3:MCF-7 whole cell lysate(40ug)

Lane4:HepG2 whole cell lysate(40ug)



Immunoprecipitation - XIAP Polyclonal Antibody

Note:

For research use only, not for use in diagnostic procedure.