

ZBP-89 (E84) polyclonal antibody

Catalog: BCP01733

Host: Rabbit

Reactivity: Human,Mouse,Rat

BackGround:

ZBP-89, also known as BFCOL1, BERF1 and ZNF 148, is a zinc finger transcription factor that is universally expressed. ZBP-89, a Kruppel-like repressor protein, is the silencer element binding factor for Vimentin. ZBP-89 has been shown to bind to GC-rich DNA elements in promoters for gastrin, ornithine decarboxylase and the cyclin-dependent kinase inhibitor p21 (also designated Cip1 or WAF1). ZBP-89 expression is induced by trans-retinoic acid or butyrate, which also induces terminal differentiation of colon cancer cells. ZBP-89 cooperates with histone acetyltransferase coactivator p300 in the regulation of p21, a cyclin-dependent kinase inhibitor whose associated gene is a target gene of p53. ZBP-89 also regulates cell proliferation, in part, through its ability to directly bind the p53 protein and retard its nuclear export. Elevated levels of ZBP-89 induce growth arrest and apoptosis in human gastrointestinal cells.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 89 kDa

Swiss-Prot:

Q9UQR1

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

ZBP-89 (E84) polyclonal antibody detects endogenous levels of ZBP-89 protein.

DATA:



Western blot (WB) analysis of ZBP-89 (E84) at 1:500 dilution


Lane1:H9C2 whole cell lysate(40ug)

Lane2:SP2/0 whole cell lysate(40ug)

Lane3:Hela whole cell lysate(40ug)

Lane4:HEK293T whole cell lysate(40ug)

Lane5:U-87MG whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of ZBP-89 (E84) pAb in paraffin-embedded human colon carcinoma tissue at 1:50, showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.