

Catenin-β (T41/S45) polyclonal antibody

Catalog: BCP01740 Host: Rabbit Reactivity: Human, Mouse

BackGround:

β-catenin is a key downstream effector in the Wnt signaling pathway. It is implicated in two major biological processes in vertebrates: early embryonic development and tumorigenesis. CK1 phosphorylates β-catenin at Ser45. This phosphorylation event primes β-catenin for subsequent phosphorylation by GSK-3 β . GSK-3 β destabilizes β-catenin by phosphorylating it at Ser33, Ser37, and Thr41. Mutations at these sites result in the stabilization of β-catenin protein levels and have been found in many tumor cell lines.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~ 90 kDa

Swiss-Prot:

P35222

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:2000~1:5000


Storage&Stability:

Store at $4\,^{\circ}$ C short term. Aliquot and store at $-20\,^{\circ}$ C long term. Avoid freeze-thaw cycles.

Specificity:

Catenin-β (T41/S45) polyclonal antibody detects endogenous levels of Catenin-β protein.

DATA:



Western blot (WB) analysis of Catenin- β (T41/S45) polyclonal antibody at 1:2000 dilution

Lane1:HEK293T whole cell lysate(30ug)

Lane2:NIH-3T3 whole cell lysate(30ug)

Note:

For research use only, not for use in diagnostic procedure.