

Complex Biotech Co., Ltd

CD1A Recombinant Protein

Catalog: 1	BCP3317	Host:	E.coli	Tag:	His-tag
------------	---------	-------	--------	------	---------

BackGround:

The CD1 multigene family encodes five forms of the CD1 T-cell surface glycoprotein in human, designated CD1A, 1B, 1C, 1D and 1E. CD1, a type 1

membrane protein, has structural similarity to the MHC class I antigen and has been shown to present lipid antigens for recognition by T lymphocytes. CD1 antigens are associated with b-2-Microglobulin and expressed on cortical thymocytes, Langerhans cells, a B cell subset and some dendritic cells. Specifically, CD1A is a marker for Langerhans cell histiocytosis (LCH) and is found on interdigitating cells. Adaptor-protein complexes and CD1-associated chaperones control CD1 trafficking, and the development and activation of CD1-restricted T cells. Constitutive endocytosis of CD1B molecules and the differential sorting of MHC class II from lysosomes separate peptide- and lipid antigen-presenting molecules during dendritic cell maturation. CD1B is also expressed in interdigitating cells. The human CD1 genes are all closely linked in a cluster mapping at chromosome 1q23.1.

Product:	Fo
PBS, 4M Urea, PH7.4	
Molecular Weight:	
~31kDa	
Swiss-Prot:	

P06126

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI

Storage&Stability: Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:

KDa	M	S
116		
66.2		
45	-	
35	-	-
25	-	
18.4		
14.4		
	*	
ST	S DAC	E Result

Note:

For research use only, not for use in diagnostic procedure.