# 

# **Complex Biotech Co., Ltd**

# **CD1B Recombinant Protein**

Catalog: BCP3318

Host: E.coli

His-tag

## **BackGround:**

The CD1 multigene family encodes five forms of the CD1 T cell surface glycoprotein in human, designated CD1A, 1B, 1C, 1D and 1E. CD1, a type 1 membrane protein, has structural similarity to the MHC class I antigen and has been shown to present lipid antigens for recognition by T lymphocytes. CD1 antigens are associated with β-2-Microglobulin and expressed on cortical thymocytes, Langerhans cells, a B cell subset and some dendritic cells. Specifically, CD1A is a marker for Langerhans cell histiocytosis (LCH) and is found on interdigitating cells. Adaptor-protein complexes and CD1-associated chaperones control CD1 trafficking, and the development and activation of CD1-restricted T cells. Constitutive endocytosis of CD1B molecules and the differential sorting of MHC class II from lysosomes separate peptide- and lipid antigen-presenting molecules during dendritic cell maturation. CD1B is also expressed in interdigitating cells. The human CD1 genes are all closely linked in a cluster mapping at chromosome 1q 23.1.

### **Product:**

PBS, 4M Urea, PH7.4

Molecular Weight:

~32kDa

**Swiss-Prot:** 

### P29016

#### **Purification&Purity:**

Tag:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

**Restriction Sites:** 

# NdeI-XhoI

Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

**Expression Vector:** 

pet-22b(+)

#### **DATA:**

KDa	$\sim$	s
116		
66.2		
45		
35		
25		
18.4		
14.4		-
	1000	

### Note:

For research use only, not for use in diagnostic procedure.