PRODUCT DATA SHEET



Complex Biotech Co., Ltd

IL-13R/CD213a1 Recombinant Protein

Catalog: BCP3370 Host: E.coli Tag: His-tag

BackGround:

The Th2 cytokine Interleukin-13 (IL-13) plays a critical role in allergen-induced airway hyper-responsiveness (AHR). Two different receptors exist for IL-13, designated IL-13Ra1 and 2. IL-13Ra1 exists as a heterodimer of IL-13Ra1 and IL-4Ra as a signaling subunit, whereas IL-13Ra2 acts as a decoy receptor for IL-13. Furthermore, TNFa or IL-4 stimulation induces IL-13Ra2 upregulation, while IL-13Ra1 is constitutively expressed. Cell surface localization of IL-13Ra2 abrogates IL-13 signaling, thus IL-13 induced translocation of the receptor from the cytoplasm provides a mechanism for negative-feedback of IL-13 signaling. IL-13Ra1 expression is predominant in B cells, monocytes and

T cells, whereas IL-13Ra2 expression is highest in glioma cells.

Product:

PBS, 4M Urea, PH7.4

Molecular Weight:

~32kDa

Swiss-Prot:

P78552

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI


Storage&Stability:

Store at $4\,^{\circ}$ C short term. Aliquot and store at $-20\,^{\circ}$ C long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:



Note:

For research use only, not for use in diagnostic procedure.