

# PRODUCT DATA SHEET

**COMPLEX**

Complex Biotech Co., Ltd

## CD117/KIT Recombinant Protein

Catalog: BCP3385

Host: E.coli

Tag: His-tag

### BackGround:

c-Kit is a member of the subfamily of receptor tyrosine kinases that includes PDGF, CSF-1, and FLT3/flk-2 receptors. It plays a critical role in activation and growth in a number of cell types including hematopoietic stem cells, mast cells, melanocytes, and germ cells. Upon binding with its stem cell factor (SCF) ligand, c-Kit undergoes dimerization/oligomerization and autophosphorylation. Activation of c-Kit results in the recruitment and tyrosine phosphorylation of downstream SH2-containing signaling components including PLC $\gamma$ , the p85 subunit of PI3 kinase, SHP2, and CrkL. Molecular lesions that impair the kinase activity of c-Kit are associated with a variety of developmental disorders, and mutations that constitutively activate c-Kit can lead to pathogenesis of mastocytosis and gastrointestinal stromal tumors. Tyr719 is located in the kinase insert region of the catalytic domain. c-Kit phosphorylated at Tyr719 binds to the p85 subunit of PI3 kinase in vitro and in vivo.

### Product:

PBS, 4M Urea, PH7.4

### Molecular Weight:

~55kDa

### Swiss-Prot:

P10721

### Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

### Restriction Sites:

NdeI-XhoI


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Expression Vector:

pet-22b(+)

### DATA:



### Note:

For research use only, not for use in diagnostic procedure.