

Complex Biotech Co., Ltd

CD159a Recombinant Protein

Tag: His-tag	ost: E.coli	BCP3416	Catalog:
--------------	-------------	---------	----------

BackGround:

The activity of natural killer (NK) cells is regulated by members of multiple receptor families that recognize class I MHC molecules, such as the killer cell inhibitory receptor/leukocyte immunoglobulin-like receptor (KIR/LIR) family and the C-type lectin superfamily. The KIR/LIR family includes p91A (also designated pp130 or PIR-B, for paired immunoglobulin-like receptor-B) and p91B (also designated PIR-A). p91A acts as an inhibitory receptor through interactions with SHP-1, whereas p91B acts as an activating receptor. CD94, NKG2 and Ly-49 are members of the C-type lectin superfamily of type II membrane glycoproteins. CD94 forms heterodimers with NKG2 isoforms on

the surface of NK cells, whereas Ly-49 isoforms form homodimers. NKG2-D, expressed on NK cells, gdT cells and CD8+ $\alpha\beta$ T cells, is a receptor for the stress inducible protein MICA, an antigen frequently expressed in epithelial tumors.

Product: PBS, 4M Urea, PH7.4 Molecular Weight: ~15kDa Swiss-Prot: P26715

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:

KDa	M	S
116		and the second se
66.2	-	
45	-	-
35	-	-
25	_	
18.4	_	-
14.4		
SD	S-PAG	E Result

Note:

For research use only, not for use in diagnostic procedure.