

PRODUCT DATA SHEET



Complex Biotech Co., Ltd

CD210 Recombinant Protein

Catalog: BCP3463

Host: E.coli

Tag: His-tag

BackGround:

The IL-10 receptor, IL-10R, is a member of the class II subgroup of the cytokine receptor family and exhibits structural similarity to the interferon receptor. IL-10R is expressed in B cells and T helper cells, as well as in LPS-induced mouse fibroblasts. Overall, mouse IL-10R and human IL-10R share 60% sequence identity at the protein level. Stimulation with IL-10 leads to phosphorylation of JAK1 and Tyk 2 tyrosine kinases. The activated kinases phosphorylate the two tyrosine residues (Tyr 446 and Tyr 496) in the cytoplasmic domain of IL-10R α . The phosphorylation of these two residues are required for proper function of IL-10R and activation of IL-10E1 signaling. IL-10R β is ubiquitously expressed and, in addition to forming the IL-10 heterodimeric receptor, it forms a heterodimeric receptor with an IL-22R subunit and an IL-28R subunit. IL-10R is constitutively expressed on human natural killer (NK) cells and the direct binding of IL-10 potentiates cytokine production by human NK cells.

Product:

PBS, 4M Urea, PH7.4

Molecular Weight:

~26kDa

Swiss-Prot:

Q13651

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:



Note:

For research use only, not for use in diagnostic procedure.