

PRODUCT DATA SHEET

COMPLEX

Complex Biotech Co., Ltd

CD218b Recombinant Protein

Catalog: BCP3471

Host: E.coli

Tag: His-tag

BackGround:

Interleukin-18 (IL-18) has been identified as a molecule that induces IFN- γ production and enhances NK cell cytotoxicity. IL-18 receptor (IL-18R) is a type I membrane protein present in lung, leukocytes, spleen, liver, thymus, prostate, small intestine, colon, placenta and heart, and absent from brain, skeletal muscle, pancreas and kidney. IL-18R is present in Hodgkin's disease cell lines, and does not bind IL-1 α or IL-1 β . The association of IL-18 to IL-18R leads to activation of NF κ B. At present two subunits of IL-18R have been characterized: IL-18R α and IL-18R β . IL-18R α has been described as the ligand-binding chain and IL-18R β as the signal-transduction chain.

Product:

PBS, 4M Urea, PH7.4

Molecular Weight:

~43kDa

Swiss-Prot:

O95256

Purification&Purity:

Transferred into competent cells and the supernatant was

purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:



Note:

For research use only, not for use in diagnostic procedure.