

PRODUCT DATA SHEET

COMPLEX

Complex Biotech Co., Ltd

CD289 Recombinant Protein

Catalog: BCP3483

Host: E.coli

Tag: His-tag

BackGround:

Members of the Toll-like receptor (TLR) family, named for the closely related Toll receptor in *Drosophila*, play a pivotal role in innate immune responses. TLRs recognize conserved motifs found in various pathogens and mediate defense responses. Triggering of the TLR pathway leads to the activation of NF- κ B and subsequent regulation of immune and inflammatory genes. The TLRs and members of the IL-1 receptor family share a conserved stretch of approximately 200 amino acids known as the Toll/Interleukin-1 receptor (TIR) domain. Upon activation, TLRs associate with a number of cytoplasmic adaptor proteins containing TIR domains, including myeloid differentiation factor 88 (MyD88), MyD88-adaptor-like/TIR-associated protein (MAL/TIRAP), Toll-receptor-associated activator of interferon (TRIF), and Toll-receptor-associated molecule (TRAM). This association leads to the recruitment and activation of IRAK1 and IRAK4, which form a complex with TRAF6 to activate TAK1 and IKK. Activation of IKK leads to the degradation of I κ B, which normally maintains NF- κ B in an inactive state by sequestering it in the cytoplasm.

TLR9 is highly expressed in macrophages, dendritic cells, and B lymphocytes, and in humans has five isoforms generated by alternative splicing. TLR9 binds to unmethylated CpG motifs present on bacterial DNA and stimulates NF- κ B via the MyD88 adaptor protein. In contrast to most TLR family members that are localized to the plasma membrane, TLR9 is an intracellular receptor localized to the ER in resting cells. Upon binding to CpG DNA, TLR9 is proteolytically processed and translocates to endo-lysosomal compartments where it binds MyD88,

initiating downstream signaling.

Product:

PBS, 4M Urea, PH7.4

Molecular Weight:

~95kDa

Swiss-Prot:

Q9NR96

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:



Note:

For research use only, not for use in diagnostic procedure.