

PRODUCT DATA SHEET

COMPLEX

Complex Biotech Co., Ltd

CD281 Recombinant Protein

Catalog: BCP3571

Host: E.coli

Tag:

His-tag;Sumo-Tag

BackGround:

Members of the Toll-like receptor (TLR) family, named for the closely related Toll receptor in *Drosophila*, play a pivotal role in innate immune responses. TLRs recognize conserved motifs found in various pathogens and mediate defense responses. Triggering of the TLR pathway leads to the activation of NF- κ B and subsequent regulation of immune and inflammatory genes. The TLRs and members of the IL-1 receptor family share a conserved stretch of approximately 200 amino acids known as the Toll/Interleukin-1 receptor (TIR) domain. Upon activation, TLRs associate with a number of cytoplasmic adaptor proteins containing TIR domains, including myeloid differentiation factor 88 (MyD88), MyD88-adaptor-like/TIR-associated protein (MAL/TIRAP), Toll-receptor-associated activator of interferon (TRIF), and Toll-receptor-associated molecule (TRAM). This association leads to the recruitment and activation of IRAK1 and IRAK4, which form a complex with TRAF6 to activate TAK1 and IKK. Activation of IKK leads to the degradation of I κ B, which normally maintains NF- κ B in an inactive state by sequestering it in the cytoplasm.

Toll-like receptor expression is highest in peripheral blood leukocytes, monocytes, macrophages, though TLR1 expression may be less restricted than other family members. TLR1 associates with TLR2 to cooperatively mediate immune responses to bacterial lipoproteins and lead to NF- κ B activation. TLR1 shows highest homology to TLR6, which shares 69% sequence identity.

Product:

0.5M Urea, PH7.4

Molecular Weight:

~43kDa

Swiss-Prot:

Q15399

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

BamHI-XhoI

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Expression Vector:

pSmart-I

DATA:



Note:

For research use only, not for use in diagnostic procedure.